PUTNAM COMPETITION 1993A

A1. The horizontal line $y = c$ intersections the curve $y = 2x - 3x^3$ in the first quadrant as in the figure. Find c such that the areas of the two shaded regions are equal.



A2. Let $(x_n)_{n\geq 0}$ be a sequence of nonzero real numbers such that

$$x_n^2 - x_{n-1}x_{n+1} = 1, \quad \text{for } n = 1, 2, 3, \ldots.$$

Prove there exists a real number a such that $x_{n+1} = ax_n - x_{n-1}$ for all $n \geq 1$.

A3. Let $\mathcal{P}_n$ be the set of subsets of $\{1, 2, \ldots, n\}$. Let $c(n, m)$ be the number of functions $f : \mathcal{P}_n \to \{1, 2, \ldots, m\}$ such that $f(A \cap B) = \min\{f(A), f(B)\}$. Prove that

$$c(n, m) = \sum_{j=1}^{m} j^n.$$

A4. Let $x_1, x_2, \ldots, x_{19}$ be positive integers each of which is less than or equal to 93. Let $y_1, y_2, \ldots, y_{93}$ be positive integers each of which is less than or equal to 19. Prove that there exists a (nonempty) sum of some x_i's equal to a sum of some y_j's.

A5. Show that

$$\int_{-100}^{-10} \left(\frac{x^2 - x}{x^3 - 3x + 1}\right)^2 dx + \int_{\frac{1}{101}}^{\frac{1}{11}} \left(\frac{x^2 - x}{x^3 - 3x + 1}\right)^2 dx + \int_{\frac{101}{100}}^{\frac{11}{10}} \left(\frac{x^2 - x}{x^3 - 3x + 1}\right)^2 dx$$

is a rational number.

A6. The infinite sequence of 2's and 3's

$$2, 3, 3, 2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 2, 3, 3, 3, 2, \ldots$$

has the property that, if one forms a second sequence that records the number of 3's between successive 2's, the result is identical to the given sequence. Show that there exists a real number r such that, for any n, the nth term of the sequence is 2 if and only if $n = 1 + \lfloor rm \rfloor$ for some nonnegative integer m. (Note: $\lfloor x \rfloor$ denotes the largest integer less than or equal to x.)